

ORIGINAL ARTICLE

# MALIGNANT MESOTHELIOMA IN THE ITALIAN REGION EMILIA-ROMAGNA: INCIDENCE AND ASBESTOS EXPOSURE UPDATE TO 2020

L. Mangone <sup>1,2</sup>, C. Storchi <sup>1</sup>, I. Bisceglia <sup>1</sup>, A. Romanelli <sup>2</sup>

<sup>1</sup> Reggio Emilia Cancer Registry, Epidemiology Unit, Azienda Unità Sanitaria Locale – IRCCS di Reggio Emilia, Reggio Emilia, Italy

<sup>2</sup> COR Emilia-Romagna, Epidemiology Unit, Azienda Unità Sanitaria Locale – IRCCS di Reggio Emilia, Reggio Emilia, Italy

## CORRESPONDING AUTHOR:

Lucia Mangone  
Reggio Emilia Cancer Registry  
COR Emilia-Romagna  
Epidemiology Unit  
Azienda Unità Sanitaria Locale – IRCCS di Reggio Emilia  
via Giovanni Amendola 2  
42122 Reggio Emilia, Italy  
E-mail: lucia.mangone@ausl.re.it  
ORCID: 0000-0003-4850-2678

Doi: 10.48286/aro.2021.20

## ABSTRACT

Malignant mesothelioma (MM) is a rare disease of great interest to the scientific community and to public health due to its high lethality and to its association with asbestos exposure.

The aim of this study, is to describe the incidence of MM in the period 1996-2020 with the related exposure to asbestos.

The data was collected by the Regional Mesothelioma Registry (ReM) which represents one of the 21 CORs (Regional Operational Centers) in Italy.

The study includes all cases of MM with a *certain* or a *probable* diagnosis (with microscopic confirmation) and *possible* cases (without microscopic confirmation).

For each case, information on sex, date of birth, tumor site, morphology, date of diagnosis, follow-up and province of residence was retrieved. We collected information on previous occupational and non-occupational exposure to asbestos, by type of sector and activity.

The exposure information is collected through the ReNaM (National Mesothelioma Registry) analytical questionnaire, administered to the patient or his closest relatives. The data collection is conducted by a regional survey network.

Were registered, between 1996 and 2020, 3013 cases of MM classified as *certain* (85.2%), *probable* (5.4%) and *possible* cases (9.4%). The greatest number of cases are recorded for the pleura (2,763) and for the peritoneum (224) in the province of Bologna, Reggio Emilia, Parma and Modena. The most affected age groups are 65-74 and 75+. Concerning exposure, 70.3% was defined *professional*, 6.0% *familiar*, 2.3% *environmental* and 1.3% *extra professional*.

As regards the provinces of residence, Bologna holds the primacy for cases of *professional* and *environmental* exposure, Reggio Emilia for *familiar* and Parma for *extra-professional* exposure. Most of the exposure to asbestos is recorded in the construction and in the railway sector, mainly in males.

## KEY WORDS

*Mesothelioma; incidence; exposure; asbestos.*

## IMPACT STATEMENT

This paper provides a 25-year overview of the incidence of malignant mesothelioma in Emilia-Romagna, a rare and highly interesting disease associated with asbestos exposure.

## INTRODUCTION

Malignant mesothelioma (MM) is a rare tumor of great scientific interest owing to its well-documented correlation with the *occupational* and/or *environmental* exposure to asbestos and the increased incidence recorded in recent years in Italy and in many other industrialized countries (1-7).

In our country, asbestos was definitively banned in April 1994 (see Law 257/92); nevertheless, the long latency time between the beginning of the exposure and the onset of the disease, the lengthening of life and the improvement of diagnostic techniques have led to an increase in the incidence of MM in recent years (8-9). MM remains a deadly cancer with a very poor prognosis, with a median of approximately 10 months from diagnosis (5, 9). In Italy, the standardized incidence rates per 100,000, recorded in 2013, are equal to 4.2 for male and 1.2 for female, whereas in individuals who were exposed to asbestos the incidence is 100-1,000 times higher. The onset generally occurs after more than 40 years of exposure to asbestos, with a median of  $48 \pm 11.4$  years (8). This pathology can also arise following modest and limited exposure to asbestos: cases have been described in workers exposed to presumably low doses and in relatives of exposed persons who took care, in a domestic environment, of the cleaning of contaminated work clothes. Cases arising from environmental exposure in residents in areas adjacent to industrial settlements where the presence or use of asbestos have also been documented (9). The ReM (Mesothelioma Registry), active since 01/01/1996, is a cancer registry specifically dedicated to the study of the incidence and etiology of MM. The objectives of the ReM, which also performs the functions of COR (Regional Operational Centers) Emilia-Romagna, are the detection of all cases of MM and the acquisition of information for a correct diagnostic definition and standardized attribution of professional or extra-work exposure to asbestos.

The aim of this study, is to report the incidence of MM in the period 1996-2020 with the related exposure to asbestos.

## MATERIALS AND METHODS

All cases of MM are detected, with pleural, pericardial, peritoneal, tunica vaginalis testis localization, arising from 1 January 1996 in subjects residing in the Emilia Romagna region at the time of diagnosis. For each registered case, in addition to the reports of the pathological investigations performed, the medical records of significant hospitalizations, carried out in public and private, regional or extra-regional health institutes, are acquired. The information on exposure, both professional and non-working, is collected through the ReNaM analytical questionnaire, administered to the patient or to his closest relatives, by the panel of occupational doctors of the Public Health Departments, members of the regional survey network. Registration for diagnostic definition and exposure attribution follow the standardized rules of the ReNaM (10). The data collection is conducted by a dedicated regional survey network that integrates all public and private Institutes and Pathological Anatomy Services operating on the regional territory, the hospital departments where patients with MM electively converge and all the Departments of Territorial Public Health. The detection network tends to acquire in *real time* the reports of new cases just diagnosed, for the early collection of the information on anamnestic exposure required directly from the patient. To verify the completeness and accuracy of incident cases, link is made with the data periodically acquired from the regional computerized archives (Mortality and Hospital Discharge Records) and information exchanges with the regional population cancer registers and the COR network.

|              | N. CASES     | DEFINITION   |
|--------------|--------------|--|
| MM certain   | 2,566        | Histology presents with characteristic morphological picture; characteristic/suggestive/absent immuno-histochemistry + diagnostic confirmation by imaging/clinical diagnosis of discharge. |
| MM probable  | 162          | Histology presents with dubious morphological picture or cytology with characteristic picture + diagnostic confirmation by imaging/clinical diagnosis of discharge.                        |
| MM possible  | 285          | Absent histology/cytology, indicative clinical and radiological data + diagnosis of MM CC discharge.   |
|              |              | DCO with wording "mesothelioma"  |
| <b>TOTAL</b> | <b>3,013</b> |  |

**Table I.** Case Distribution by diagnostic definition, cases from 1996 to 2020 (updated to 12/31/2020).

| YEAR                   | N. CASES     |            |             |           | TOTAL        |
|------------------------|--------------|------------|-------------|-----------|--------------|
|                        | SITE         |            |             |           |              |
|                        | PLEURA       | PERITONEUM | PERICARDIUM | TESTIS    |              |
| 1996                   | 63           | 8          | -           | 2         | 73           |
| 1997                   | 70           | 7          | 3           | -         | 80           |
| 1998                   | 77           | 4          | 1           | 1         | 83           |
| 1999                   | 67           | 6          | -           | -         | 73           |
| 2000                   | 76           | 9          | -           | 1         | 86           |
| 2001                   | 88           | 6          | -           | 2         | 96           |
| 2002                   | 98           | 15         | -           | 1         | 114          |
| 2003                   | 97           | 6          | 1           | 1         | 105          |
| 2004                   | 110          | 8          | 2           | -         | 120          |
| 2005                   | 107          | 10         | -           | 2         | 119          |
| 2006                   | 100          | 7          | -           | -         | 107          |
| 2007                   | 101          | 14         | -           | -         | 115          |
| 2008                   | 122          | 9          | -           | 1         | 132          |
| 2009                   | 111          | 11         | -           | -         | 122          |
| 2010                   | 117          | 12         | 1           | -         | 130          |
| 2011                   | 144          | 10         | -           | 1         | 155          |
| 2012                   | 142          | 10         | 1           | 2         | 155          |
| 2013                   | 147          | 5          | -           | 1         | 153          |
| 2014                   | 122          | 10         | -           | 1         | 133          |
| 2015                   | 141          | 10         | -           | -         | 151          |
| 2016                   | 150          | 10         | -           | -         | 160          |
| 2017                   | 146          | 11         | -           | 1         | 158          |
| 2018                   | 138          | 14         | -           | -         | 152          |
| 2019                   | 132          | 8          | -           | -         | 140          |
| 2020*                  | 97           | 4          | -           | -         | 101          |
| <b>TOTAL 1996-2020</b> | <b>2,763</b> | <b>224</b> | <b>9</b>    | <b>17</b> | <b>3,013</b> |

**Table II.** Distribution of cases by site and year of diagnosis.

\* The incidence is not complete.

## RESULTS

In Emilia Romagna 3013 MM were registered between 1996 and 2020: the distribution by type of diagnostic definition is shown in **table I**. Most of the cases (2,566) were classified as *certain* (85.2%), 162 *probable* (5.4%) and 285 (9.4%) *possible* cases and 6 cases with only the death certificate (**table I**). The distributions by year of diagnosis and localization is shown in **table II**. The cases go from less than 100 per year in the first six years of registration to an average of about 110 in 2003-2007, 130 in 2008-2010 and then reaching around 150 starting from 2011.

In the period 1996-2020 the greatest number of cases are recorded for the pleura (2,763) and for the peritoneum (224), on average 10 cases per year. Tumours in the testis are rare (17), about 1 per year. Those of the pericardium are rare.

Table 3 shows the distribution by year and residence. The provinces in which the majority of cases are recorded are: Bologna (660), followed by Reggio Emilia (463), Parma (355) and Modena (329).

An overview of the distribution of MM in the Emilia-Romagna region referring to the period 2014-2018 (the most recent complete five-year period) is shown in **figure 1**. Compared with the national data for 2013 (males 4.2 and females 1.2), Emilia

Romagna has a slightly lower rate in males (3.9) and higher in females (1.4).


Compared with the regional data of Emilia Romagna for the period 2014-2018 (3.8 in males and 1.1 in females), it is the province of Reggio Emilia that shows the highest rates in males (5.7) and females (2.3). Following, in males, the provinces of Piacenza (4.9) and Ravenna (4.6); while in the females Piacenza (1.4), Ferrara (1.3) and Parma (1.2).

As for age (**table IV**), there is a strong gradient for pleural MM in both sexes. Most of the cases are concentrated in the age of 65+. For the peritoneum, most cases, regardless of age, are recorded at the age of 65+; while there are no differences for testis and pericardium.

The incidence trend (**figure 2**) shows a slight increase in males but not females.

Of the 3,013 cases of MM, information on exposure to asbestos was collected in 2,605 cases, 169 not defined and 239 not classifiable. Overall, considering all those exposed to asbestos, in our region it was found that 80% of respondents were exposed (87% males and 60% females) (**figure 3**).

The distribution of type of exposure is shown in **table V**: in 70.3% it was defined *professional*, 6% *familiar*, 2.3% *environmental* and 1.3% *extra profes-*



**Figure 1.** ReM cases 1995-2020. Distribution of cases and TSD by province of residence.




Figure 2. Standardized incidence rates on the Italian population x 100,000. Years 1996-2018.




Figure 3. Distribution by exposure to asbestos by gender in the Emilia-Romagna region.

sional. Occupational exposure is much higher in males (83%) while the family exposure is higher in females (20.6%).

As regards the provinces of residence, Bologna holds the primacy for cases of professional (400) and environmental (19) exposure, Reggio Emilia for familiar (38) and Parma for extra-professional exposure (7) (table VI).

Table VII shows the working sector involved: most of the exposure to asbestos is recorded in the construction sector and in the railway sector, mainly

in males. Women, in general, are less affected, but most of the cases are recorded in the sectors of sugar refineries and in production of cement.

## DISCUSSION

Malignant Mesothelioma confirms its characteristics of rare tumor with an increase in the incidence in both genders, recorded up to 1996 and with the first signs of a steady trend in the following years.

| YEAR         | N. CASES   |            |            |            |            |            |            |            |            |              |
|--------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|--------------|
|              | PC         | PR         | RE         | MO         | BO         | FE         | RA         | FC         | RN         | R E-R        |
| 1996         | 5          | 12         | 13         | 8          | 18         | 7          | 7          | 1          | 2          | 73           |
| 1997         | 9          | 9          | 10         | 3          | 24         | 7          | 5          | 7          | 6          | 80           |
| 1998         | 8          | 8          | 13         | 11         | 20         | 7          | 7          | 6          | 3          | 83           |
| 1999         | 7          | 6          | 10         | 6          | 14         | 9          | 7          | 8          | 6          | 73           |
| 2000         | 7          | 10         | 10         | 8          | 17         | 13         | 8          | 11         | 2          | 86           |
| 2001         | 9          | 13         | 12         | 8          | 22         | 17         | 5          | 5          | 5          | 96           |
| 2002         | 8          | 7          | 16         | 12         | 37         | 13         | 11         | 8          | 2          | 114          |
| 2003         | 11         | 7          | 16         | 11         | 24         | 10         | 12         | 7          | 7          | 105          |
| 2004         | 8          | 16         | 18         | 10         | 24         | 10         | 16         | 12         | 6          | 120          |
| 2005         | 13         | 24         | 16         | 12         | 22         | 13         | 9          | 7          | 3          | 119          |
| 2006         | 15         | 15         | 10         | 10         | 29         | 10         | 11         | 5          | 2          | 107          |
| 2007         | 4          | 19         | 22         | 15         | 28         | 6          | 13         | 6          | 2          | 115          |
| 2008         | 9          | 14         | 13         | 20         | 23         | 18         | 13         | 12         | 10         | 132          |
| 2009         | 7          | 16         | 10         | 13         | 26         | 16         | 19         | 10         | 5          | 122          |
| 2010         | 12         | 16         | 19         | 11         | 26         | 9          | 16         | 14         | 7          | 130          |
| 2011         | 14         | 14         | 22         | 22         | 31         | 15         | 18         | 10         | 9          | 155          |
| 2012         | 15         | 21         | 29         | 13         | 34         | 16         | 12         | 7          | 8          | 155          |
| 2013         | 15         | 16         | 25         | 11         | 35         | 18         | 18         | 11         | 4          | 153          |
| 2014         | 12         | 16         | 25         | 19         | 28         | 8          | 16         | 5          | 4          | 133          |
| 2015         | 12         | 13         | 20         | 20         | 33         | 23         | 14         | 8          | 8          | 151          |
| 2016         | 11         | 17         | 27         | 22         | 33         | 12         | 11         | 13         | 14         | 160          |
| 2017         | 17         | 13         | 27         | 20         | 35         | 14         | 15         | 11         | 6          | 158          |
| 2018         | 12         | 14         | 36         | 17         | 28         | 11         | 17         | 7          | 10         | 152          |
| 2019         | 13         | 22         | 20         | 15         | 24         | 11         | 19         | 10         | 6          | 140          |
| 2020         | 5          | 17         | 25         | 11         | 25         | 4          | 7          | 5          | 2          | 101          |
| <b>TOTAL</b> | <b>258</b> | <b>355</b> | <b>463</b> | <b>329</b> | <b>660</b> | <b>297</b> | <b>306</b> | <b>206</b> | <b>139</b> | <b>3,013</b> |

**Table III.** Distribution of cases by province of residence and year of diagnosis.

| AGE          | MALE         |            |             |           | FEMALE     |            |             | TOTAL        |
|--------------|--------------|------------|-------------|-----------|------------|------------|-------------|--------------|
|              | N. CASES     |            |             |           | N. CASES   |            |             |              |
|              | Pleura       | Peritoneum | Pericardium | Testis    | Pleura     | Peritoneum | Pericardium |              |
| < 45         | 18           | 9          | -           | 4         | 11         | 5          | -           | 47           |
| 45-54        | 88           | 10         | -           | 3         | 36         | 10         | 1           | 148          |
| 55-64        | 336          | 22         | -           | 2         | 94         | 21         | 2           | 477          |
| 65-74        | 683          | 44         | 4           | 3         | 224        | 29         | 1           | 988          |
| 75+          | 904          | 43         | 1           | 5         | 369        | 31         | -           | 1,353        |
| <b>TOTAL</b> | <b>2,029</b> | <b>128</b> | <b>5</b>    | <b>17</b> | <b>734</b> | <b>96</b>  | <b>4</b>    | <b>3,013</b> |

**Table IV.** Distribution of cases by sex, site and age at diagnosis.

| TYPE OF EXPOSURE              | MALE         |              | FEMALE     |              | TOTAL        |              |
|-------------------------------|--------------|--------------|------------|--------------|--------------|--------------|
|                               | N. CASES     | %            | N. CASES   | %            | N. CASES     | %            |
| Professional                  | 1,607        | 83.5         | 225        | 33.1         | 1,832        | 70.3         |
| Familiar                      | 15           | 0.8          | 140        | 20.6         | 155          | 6.0          |
| Environmental                 | 29           | 1.5          | 30         | 4.4          | 59           | 2.3          |
| Extra Professional            | 20           | 1.0          | 15         | 2.2          | 35           | 1.3          |
| Improbable                    | 63           | 3.3          | 87         | 12.8         | 150          | 5.8          |
| Unknown                       | 191          | 9.9          | 183        | 26.9         | 374          | 14.3         |
| <b>TOTAL OF DEFINED CASES</b> | <b>1,925</b> | <b>100.0</b> | <b>680</b> | <b>100.0</b> | <b>2,605</b> | <b>100.0</b> |

**Table V.** Distribution of malignant mesotheliomas by type of exposure and sex.

| TYPE OF EXPOSURE   | N. CASES   |            |            |            |            |            |            |            |            |              |
|--------------------|------------|------------|------------|------------|------------|------------|------------|------------|------------|--------------|
|                    | PC         | PR         | RE         | MO         | BO         | FE         | RA         | FC         | RN         | R E-R        |
| Professional       | 147        | 219        | 314        | 163        | 400        | 177        | 215        | 119        | 78         | 1,832        |
| Familiar           | 11         | 20         | 38         | 7          | 32         | 19         | 8          | 9          | 11         | 155          |
| Environmental      | 4          | 13         | 7          | 5          | 19         | 4          | 3          | 4          | -          | 59           |
| Extra-professional | 5          | 7          | 1          | 4          | 6          | 2          | 4          | 4          | 2          | 35           |
| Improbable         | 12         | 20         | 8          | 15         | 41         | 14         | 24         | 8          | 8          | 150          |
| Unknown            | 40         | 48         | 26         | 40         | 93         | 33         | 40         | 34         | 20         | 374          |
| To be defined      | 6          | 10         | 60         | 42         | 33         | 4          | 6          | 7          | 1          | 169          |
| Not classifiable   | 33         | 18         | 9          | 53         | 36         | 44         | 6          | 21         | 19         | 239          |
| <b>TOTAL</b>       | <b>258</b> | <b>355</b> | <b>463</b> | <b>329</b> | <b>660</b> | <b>297</b> | <b>306</b> | <b>206</b> | <b>139</b> | <b>3,013</b> |

**Table VI.** Distribution of cases by type of exposure and province of residence.

It should be noted that the ReM data relating to 2019 and 2020, which show a significant decrease, of -8 % and -34% respectively, compared with the incidence rates recorded in 2018, are probably due to the detection deficit caused by the current viral pandemic, which makes it necessary to adopt ad hoc in-depth studies currently in progress at the Registry with the involvement of the ReNaM CORs. In view of its almost total lethality, however, this disease still assumes social relevance with an impact greater than that of fatal injuries. The INAIL fatal injuries data reported in 2015-2019 continue to demonstrate a lower occurrence, 610 vs 761, compared to the incidence of MM, recorded by the ReM in the same period (see INAIL Annual Report, June 2020). Since the disease is almost always associated with exposures, even modest ones, to asbestos, each new case must be considered a "sentinel event" of previous exposures and carefully evaluated (7, 10-20). Based on these considerations, the primary objective of the ReM is certainly the completeness of

the data and the accuracy of the information collected. These aims seem to have been achieved thanks to the capillary regional detection network which also allows a good recording of MM with extra pleural localization. The diagnostic quality can be considered of a good level: 90.5% of cases are accompanied by cyto-histological confirmation owing to the widespread practice in the regional health services of performing biopsy with minimally invasive techniques, which allow examining also elderly patients and/or patients with reduced "performance status". The involvement of Prevention Services is certainly important because it guarantees a correct anamnestic reconstruction of exposure and the drafting of good quality certifications for INAIL to guarantee the patient and his family members the recognition of the privileged public protection provided for technopathies. In all our cases, these are 249 subjects out of 2,081 with ascertained exposure to asbestos (12.0%), for which it is not possible, under current legislation, to access privileged forms



| PRODUCTION SECTOR                                   | MALE         |              | FEMALE     |              | TOTAL        |              |
|---|--------------|--------------|------------|--------------|--------------|--------------|
|   | N. CASES     | %            | N. CASES   | %            | N. CASES     | %            |
| Constructions                                       | 269          | 16.7         | 1          | 0.4          | 270          | 14.7         |
| Constructions/repair railway rolling stocks         | 186          | 11.6         | 3          | 1.3          | 189          | 10.3         |
| Engineering industry                                | 151          | 9.4          | 12         | 5.4          | 163          | 8.9          |
| Sugar refineries/other food industries              | 112          | 7.0          | 37         | 16.4         | 149          | 8.1          |
| Production of cement/asbestos products              | 93           | 5.8          | 32         | 14.2         | 125          | 6.8          |
| Production of chemical/plastic material             | 98           | 6.1          | 6          | 2.7          | 104          | 5.7          |
| Building completion works                           | 80           | 5.0          | 1          | 0.4          | 81           | 4.4          |
| Glass/ceramic/rubber manufacturing                  | 56           | 3.5          | 20         | 8.9          | 76           | 4.2          |
| Transportation                                      | 72           | 4.5          | 3          | 1.3          | 75           | 4.1          |
| Production/repair vehicles<br>(no trains and ships) | 65           | 4.0          | 3          | 1.3          | 68           | 3.7          |
| Manufacturing/processing of metallic products       | 60           | 3.7          | 3          | 1.3          | 63           | 3.5          |
| Textile industry                                    | 35           | 2.2          | 15         | 6.7          | 50           | 2.7          |
| Trade   | 37           | 2.3          | 8          | 3.6          | 45           | 2.5          |
| Production electricity, gas, water                  | 39           | 2.4          | -          | -            | 39           | 2.1          |
| Social services/recreational activities/healthcare  | 19           | 1.2          | 17         | 7.6          | 36           | 2.0          |
| National defense                                    | 35           | 2.2          | 1          | 0.5          | 36           | 2.0          |
| Agriculture/animal breeding                         | 23           | 1.4          | 12         | 5.4          | 35           | 1.9          |
| Metallurgical industry                              | 27           | 1.7          | 4          | 1.8          | 31           | 1.7          |
| Other manufacturing industries                      | 27           | 1.7          | 3          | 1.3          | 30           | 1.6          |
| Other   | 123          | 7.6          | 44         | 19.6         | 167          | 9.1          |
| <b>TOTAL</b>  | <b>1,607</b> | <b>100.0</b> | <b>225</b> | <b>100.0</b> | <b>1,832</b> | <b>100.0</b> |

**Table VII.** Distribution of occupational exposure to asbestos by main sector of economic activity and sex.

of protection for damage from work.

The extension of the asbestos victims' fund to people suffering from MM due to exposure to "non-professional" asbestos tends to overcome this situation with the provision of an indemnity, albeit *una tantum*; this was introduced on an experimental basis for the three-year period 2015-2017 by the established law 2015 (see L 190/2014 and DIM 04/09/2015), validated for the following three-year period 2018-2020 and increased pursuant to art. 11 quinquies, Law 8/2020.

Significantly, the MM recorded in Emilia-Romagna show a high proportion of cases with extra pleural site: the pleura / extra pleural ratio recorded by the ReM was equal to 11.1:1, compared to 13.4:1 and 13,3:1 recorded by CORs in Italy (8, 11) and to some international reports (12, 14) which probably underestimate the data of MM with extra pleural site. Certainly, the articulation of the ReM detection network favours the exhaustiveness of the information collection both from the clinical departments, pulmonology and thoracic surgery mainly,

where the MM with pleural site electively flow, and from those where cases with extra pleural site are treated: general surgery, gynaecology, cardiac surgery, urology and andrology. On the other hand, research carried out in the ISPESL/ReNaM field had highlighted some difficulties, especially in some CORs, in the systematic detection of MM with an extra pleural site and had identified possible ways to implement this detection (11).

As regards the age at diagnosis, the average was  $71.9 \pm 10.7$  years; it is noteworthy that 77.7% of subjects were <sup>3</sup> 65 years of age at the time of diagnosis compared to 72.0% recorded in Italy (8). The data could be correlated to a greater tendency, in our region, to perform biopsy sampling even in older subjects, thanks to the good diffusion of minimally invasive practices (e.g. video-thoracoscopy) compared to more aggressive traditional methods. The regional annual incidence rates per 100,000, standardized for the 2000 Italian population, show an increasing trend. The years with a higher incidence were 2011, 2012 and 2016 for male (4.4)



and 2013 and 2018 for female (1.4) (**figure 2**). The 2014-2018 regional average rates (3.8 M and 1.1 W) are lower than those recorded by the ReNaM in 2013 (4.2 M and 1.2 W).

The 2014-2018 TIS show data that cannot be readily interpreted for Piacenza and Ferrara, whereas for Reggio Emilia they are mainly correlated to the significant spread in the past of companies engaged in the production of asbestos-cement products and the construction/repair of railway rolling stock. In particular, the high value for female is certainly due to the use, peculiar in this province, of female engagement in the manufacturing of "special pieces" in cement/asbestos. The analysis of exposure to asbestos for the 2,605 cases already investigated, highlighted exposure in 79.9% of cases, whereas for the remaining 20.1% no information was found concerning exposure to asbestos, which defines the class "unlikely/unknown" asbestos exposure.

This result, rather than indicating a real absence of previous exposures, even remote and episodic, is likely to be ascribed to the difficulty of recording exhaustive anamnestic, professional or extra-professional exhibition information, concerning situations that could have occurred even a few decades earlier the onset of the disease. These difficulties, more relevant for the female gender, are also linked to the reduced median survival of the MM which does not always allow to detect good quality information from the patient voice.

In most of the subjects exposed to asbestos, the origin of the exposure was traced to professional activities (88.0%), whereas exposure due to cohabitation with professionally exposed subjects or to extra-work activities occurred in 9.1% of cases. Also, among the subjects exposed to asbestos in our Region, the rate of those who developed MM because "they lived near production sites that worked or used asbestos (or materials containing asbestos)" or they have frequented environments with the presence of asbestos for non-professional reasons", the so-called environmental exposure to asbestos (see LL.GG. ReNAM) (10), is equal to 2.9%. This fraction is lower than that recorded by ReNaM in Italy, which is equal to 4.4%, and then that recorded in some Italian municipalities, subject in the past to significant environmental contamination by asbestos.

In the Emilia-Romagna Region, the production

sectors most involved in the onset of MM were: building constructions (subjects distributed evenly throughout the region); construction/repair of railway rolling stock (cases mostly resident in the provinces of Bologna and Reggio Emilia); metal-working industry, sugar refineries/other food industries (118 of 149 cases, residing in the provinces of BO, FE, RA, PR, FC), production of asbestos cement products (98 of the 125 cases residing in the province of RE). National ReNaM data, on the other hand, indicate among the most involved sectors, in addition to construction (15.5%) and the engineering industry (8.6%), the textile industry (6.4) and shipbuilding (6.1%).

## CONCLUSIONS

The study shows that MM is a rare but still highly lethal pathology. The early diagnosis of the neoplasm, although not able to influence the prognosis, allows to obtain more precise information directly from the patient on previous exposure to asbestos and to ensure, in due cases, the right compensation to those who have paid too high a price during his job duties.

## ACKNOWLEDGMENTS

The collection, archiving and definition of malignant MM cases incident throughout the Emilia-Romagna Region was possible, with an acceptable cost/benefit ratio, only through the effective collaboration and careful development of the regional detection network. which has over 140 formally designated Referents<sup>1</sup>, including: pathologist specialists, hygienists and occupational doctors from the Departments of Public Health, pulmonologists, general surgeons, gynaecologists, urologists, oncologists, but also internists and cardiologists that the good collaboration established can guarantee an increasingly adequate knowledge of this fearful pathology.

## CONFLICT OF INTERESTS

The authors have declared no conflict of interests.

## REFERENCES

1. Novello S, Pinto C, Torri V, et al. The Third Italian Consensus Conference for Malignant Pleural Mesothelioma: State of the art and recommendations. *Crit Rev Oncol Hematol* 2016;104:9-20.
2. Ministero della Salute. Piano Nazionale Amianto: Linee di intervento per un'azione coordinata delle amministrazioni statali e territoriali; Roma, marzo 2013.
3. Hughes S. Relazione sulle minacce per la salute sul luogo di lavoro legate all'amianto e le prospettive di eliminazione di tutto l'amianto esistente; Parlamento Europeo, Doc di seduta A7- 0025/2013.
4. Marinaccio A, Binazzi A, Di Marzio D, et al. Pleural malignant mesothelioma epidemic. Incidence, modalities of asbestos exposure and occupation involved from the Italian National Register. *Int J Cancer* 2012;130(9):2146-54.
5. Alessi M, Amadori D, Amunni G, et al. Stato dell'arte e prospettive in materia di contrasto alle patologie asbesto-correlate; Quaderni del Ministero della Salute, n° 15, maggio-giugno 2012.
6. Delgermaa V, Takahashi K, Park EK, Vinh Le G, Hara T, Sorahan T. Global mesothelioma deaths reported to the World Health Organization between 1994 and 2008. *Bull World Health Organ* 2011;89:716-24.
7. Bertazzi PA. Descriptive epidemiology of malignant mesothelioma. *Med Lav* 2005;7(4):287-303.
8. Marinaccio A, Binazzi A, Branchi C, et al. Sesto Rapporto – Il Registro Nazionale dei Mesoteliomi. INAIL, Milan 2018.
9. Magnani C, Bianchi C, Chellini E, et al. III Consensus Conference on Malignant Mesothelioma of the Pleura. *Epidemiology, Public Health and Occupational Medicine related issues. Med Lav* 2015;106(5):325-32.
10. Nesti M, Adamoli S, Ammirabile F, et al. Linee Guida per la rilevazione e la definizione dei casi di mesotelioma maligno e la trasmissione delle informazioni all'ISPESL da parte dei Centri Operativi Regionali. II Edizione. Roma, maggio 2004.
11. Romanelli A, Marinaccio A, Mirabelli D, et al.. Progetto di ricerca ISPESL B/45/DML/03, I mesoteliomi maligni a localizzazione extrapleurica, 2005.
12. Robinson BW, Musk AW, Lake RA. Malignant mesothelioma. *Lancet* 2005;366:397-408.
13. Chiappino G, Mensi C, Riboldi L, Rivolta G. Il rischio amianto nel settore tessile: indicazioni dal Registro Mesoteliomi Lombardia e definitiva conferma. *Med Lav* 2003;94(6):521-30.
14. Sugarbaker PH, Welch LS, Mohamed F, Glehen O. A review of peritoneal mesothelioma at the Washington Cancer Institute. *Surg Oncol Clin N Am* 2003;12(3):605-21.
15. Britton M. The epidemiology of mesothelioma. *Semin Oncol* 2002;29(1):51-61.
16. Huncharek M. Non-asbestos related diffuse malignant mesothelioma. *Tumori* 2002;88:1-9.
17. Mangone L, Romanelli A, Campari C, Candela S. Il mesotelioma maligno in Emilia-Romagna: incidenza ed esposizione ad amianto. *Epid Prev* 2002;26(3):124-9.
18. Peto J, Decarli A, La Vecchia C, Levi F, Negri E. The European mesothelioma epidemic. *Br J Cancer* 1999;79(34):666-72.
19. Boffetta P. Health effects of asbestos exposure in humans: a quantitative assessment. *Med Lav* 1998;89(6):471-80.
20. Spirtas R, Heineman EF, Bernstein L, et al. Malignant mesothelioma: attributable risk of asbestos exposure. *Occup Environ Med* 1994;51:804-11.